



F.No. 455755-F/9(4)2024

भारत सरकार

Government of India

कृषि एवं किसान कल्याण मंत्रालय

Ministry of Agriculture & Farmers Welfare

कृषि एवं किसान कल्याण विभाग

Department of Agriculture & Farmers Welfare

वनस्पति संरक्षण, संगरोध एवं संग्रह निदेशालय

Directorate of Plant Protection, Quarantine & Storage

केंद्रीय कीटनाशी बोर्ड एवं पंजीकरण समिति

Central Insecticides Board and Registration Committee

एन. एच. 4, फरीदाबाद (हरियाणा)-121001

N.H. IV, FARIDABAD (HARYANA)-121001

CERTIFICATE OF REGISTRATION UNDER SECTION 9(4) OF THE INSECTICIDES ACT, 1968.

Certified that the Insecticide Acetamiprid 20% SP has been registered under section 9(4) of the Act in the name of the Person/Undertaking whose particulars are specified below:-

1. Name of the person/Undertaking : M/sSAHAJ AGROCHEM INDUSTRIES,SF 37, Earth Alpha, Vasna Road, Vadodara- 390007 (Gujarat),Gujarat,Vadodara,Vadodara,390007
2. Address of the manufacturing premises : 19/D, Raj Industrial Park, Karchiya, Savli, Vadodara - 391310 (Gujarat) .
3. Registration Number : CIR-346880/2025-Acetamiprid (SP) (461)-1443
4. Name of the Insecticide : Acetamiprid 20% SP

5. CONDITIONS :

- i. The insecticide shall be manufactured indigenously.
- ii. The registration is subject to the strict compliance of various provisions of the Insecticides Act,1968 as amended from time to time and Rules, bye-laws framed and notifications issued thereunder and as amended from time to time.
- iii. The registration certificate is further subject to such conditions which may be varied and specified from time to time by the Registration Committee under section 9(3c).
- iv. The registration Certificate holder shall strictly comply with the condition and amendments thereof set out in this certificate of registration (CR) including label and leaflets. Non-compliance of the condition set out herein before and hereinafter will entail action under section 17 of the Act.
- v. The insecticide shall have the composition (kind, name and percentage of the ingredients) as given below :-

S.No	Component	Component Desc.	Content (% w/w)
1	Acetamiprid a.i.		20.00 % w/w
2	Colouring agent	(Brilliant Blue)	0.05 % w/w
3	Sodium Sulphate & Sodium alkylate Mixture		% w/w
4	Blend of Sodium Dodecyl Benzene Sulfonate		4.50 %
5	Sodium Citrate, Dihydrate		30.00 % w/w
6	Calcium Hydrogen Phosphate Dihydrate		0.10 % w/w

- vi. The insecticides shall contain the maximum impurities as quantified/identified and submitted to the Registration Committee.
- vii. The Product shall confirm to the specification submitted by you and also to the IS vide No. and amendment thereof as and when the same are formulated and published.
- viii. A sample quantity of the insecticide being registered along with a small quantity of reference analytical standard should be sent to the Director, Central Insecticides Laboratory, Directorate of Plant Protection, Quarantine & Storage, N.H. IV, Faridabad, as and when required, for verification.
- ix. A copy, each of the approved label and leaflet is enclosed. No change, addition, alteration, modification or deletion with respect to the inscriptions on the labels/leaflets shall be done without the prior approval of the Registration Committee.
- x. The labels and leaflets shall be printed by using letters that are BOLD enough for a man of ordinary/normal vision to read them without any external help.
- xi. The Registration Committee does not itself responsible for the use of trade name by you. The use of the trade name shall be regulated as per the existing laws on the subject.
- xii. The license should be granted subject to the conditions that the licensee shall comply with the provisions of the Act and the Rules made thereunder and the conditions of registration for the time being in force.
- xiii. No license to manufacture an insecticide shall be granted unless the licensing officer is satisfied that necessary plant and machinery, safety devices, first-aid facilities, quality control measures, the requirement laid down under Chapter VIII of the Rules, etc. exist in the premises where the insecticide is proposed to be manufactured.
- xiv. The shelf-life of the insecticide shall be 2 year(s) .
- xv. Inspection of the manufacturing unit should be undertaken to collect in-process samples of the insecticides. The samples should be analyzed to verify the claims made by the licensee relating to chemical parameters, and the report thereof should be submitted to the Registration Secretariat within a period of six months.
- xvi. No export should take place in contravention to the provisions of the Rotterdam Convention on prior informed consent procedure for certain hazardous chemicals and pesticides in international trade.
- xvii. If a pesticide is banned or severely restricted in India, before exporting such pesticide, permission from Designated National Authority for Pesticide of the Country under Rotterdam Convention may be obtained.
- xviii. Health records of Industrial workers may be maintained in Appendix E (conditions) of Form-II as prescribed in the Insecticides Rules 1971. In case any untoward/adverse effect is noticed, then the same may be reported to Registration Secretariat by the Licensing Officer.
- xix. The registrant shall have to commence actual production of the pesticide within three years from the date of issue of registration certificate and is required to produce a certificate from the concerned State/UT Governments as a proof of production, failing which the certificate of registration shall automatically lapse.
- xx. In case of export, the packaging shall be as per the requirement of the importing country and conforming to IMDG guidelines.
- xxi. The registrant has to submit detail of Import/Export or indigenous manufacturing (as the case may be) of this pesticide month wise mandatory to the Secretariat of CIB&RC as the case maybe. In case non compliance of this condition is observed this CR shall be canceled immediately without any Notice.
- xxii. The product is registered for domestic use as well as for Export.
- xxiii. The product is registered for domestic as well as for export and in case of export primary packaging shall be as per the requirement of importing country.
- xxiv. The registrant is required to use the label and leaflets as per GSR 355(E) dated 04.06.2020 and applicable

Insecticides Rules. The Issuance of the Certificate of Registration is strictly subject to the outcome of the W.P(C) No. 4136/2020 and W.P(C) No. 4137/2020.

Specific Conditions

The product shall be packed in multilayer laminated pouches comprising of polyester 12 micron/Al-foil 9 micron/LDPE 0.062 mm only. The capacity of the pouch shall be 20g., 50g., 100g., 250g., 500g., and 1.00 Kg. only. These shall be individually inserted in card board carton which shall be further packed in CFB boxes. The capacity of CFB boxes containing primary packaging upto 500g. size shall be 2.00 Kgs. and containing primary packaging for 1.00 Kg. shall be 5.00 Kg. only. No other packaging system shall be used without the approval of the Registration Committee.

Dated: 24/02/2025

Copy to: The Director of Agriculture ,Gujarat



**Dr. Archana Sinha
Secretary
Central Insecticides Board
and Registration Committee**

Note :-

The Labels and leaflets have been generated through newly created data bank. Notwithstanding diligence exercised in creation of the data bank, the possibility of errors creeping into labels and leaflets cannot be ruled out. The same is subjected to necessary corrections.

Signature valid

Digitally signed by Archana Sinha
Date: 2025.02.28 16:17:18
Location: CIB&RC,NH4,CGO
Complex, Faridabad.
Designation: Secretary (CIB&RC)



Label Details

Chemical Composition

S.No	Ingredient	Description	Weight
1	Acetamiprid a.i.		20.00 % w/w
2	Colouring agent	(Brilliant Blue)	0.05 % w/w
3	Sodium Sulphate & Sodium alkylate Mixture		% w/w
4	Blend of Sodium Dodecyl Benzene Sulfonate		4.50 %
5	Sodium Citrate, Dihydrate		30.00 % w/w
6	Calcium Hydrogen Phosphate Dihydrate		0.10 % w/w
7	Lactose Monohydrate		Q.S %
Total			100 % w/w

Direction Of Use

Please read enclose leaflet before use.

Precautions

1. Keep away from foodstuffs, empty foodstuff containers and animals food. 2. Avoid contact with mouth, eyes and

CIR-346880/2025-Acetamiprid (SP) (461)-


1443

Acetamiprid 20% SP

(Insecticide)

ऐसिडामिप्रिड 20 प्रतिशत एस पी

(कीटनाशक)



बच्चों की पहुँच से दूर रखें

Keep out of reach of children.

सूखी और ठण्डी जगह पर आग से दूर रखें

Keep in cool,dry place away from heat and open flame.

Regn.No रजि. नं. _____
Mfg.Lic.No मैनयु ला. नं. _____
Batch No. बैच नं. _____
Mfg Date मैनयु, ता. _____

रासायनिक संरचना

क्रमांक	घटक	विवरण	वजन
1	ऐसिडामिप्रिड सक्रि. तत्व		20.00 % भार/ भार
2	रंग लाने वाले घटक	(ब्रिलीयन्ट ब्लू)	0.05 % भार/ भार
3	सोडियम सल्फनेट एण्ड सोडियम एल्केलाईट		%
4	सोडियम डोडिसाइल बेन्जीन सल्फोनेट		4.50 % भार/ भार
5	सोडियम साइट्रेट, डाईहाइड्रेट		30.00 % भार/ भार
6	कैल्शियम हाइड्रोजन फॉस्फेट डाईहाइड्रेट		0.10 % भार/ भार
7	लैक्टोज मोनोहाइड्रेट		पर्याप्त मात्रा %
			कुल 100 % w/w

उपयोग के लिए दिशा-निर्देश

उपयोग से पहले संलग्न लीफलेट पढ़ें ।

skin. 3. Avoid inhalation the spray mist. Spray in the direction of wind. 4 . Wash thoroughly the contaminated clothes and parts of the body after spraying. 5. Do not smoke, drink, eat and chew anything while spraying. 6. Wear full protective clothing while mixing and spraying.

Antidote

No specific antidote is known. Treat symptomatically.

First Aid

Please see enclosed Leaflet.

Symptoms

Please see enclosed Leaflet.

Recommendations

Acetamiprid 20% SP is a soluble powder formulation containing 20% of the active ingredient Acetamiprid. Acetamiprid 20% SP is a highly effective systemic insecticide for the control of aphids, jasssids and whiteflies in cotton crop; Aphid in Cabbage; Aphid in Okra.

Manufactured By

M/s SAHAJ AGROCHEM INDUSTRIES

SF 37, Earth Alpha, Vasna Road, Vadodara- 390007 (Gujarat)

Manufacturer Premises Address

Exp. Date	खत्म होने की	_____
	तारीख	
Net Content / Volume	शुद् भार,	_____
	आयतन	
Max. Retail Price	अधिकतम	_____
	खुदरा मूल्य	

प्रयोगकर्ताओं के लिए सावधानियां

1. खाध सामग्री,खाध सामग्री के खाली बर्तन और पशुओं के चारे से दूर रखें।
2. मुँह, त्वचा और आंखों के सम्पर्क से बचायें।
3. छिड़काव की वाष्प को सांस द्वारा अन्दर जाने से बचायें। हवा की दिशा में छिड़काव करें।
4. छिड़काव के बाद दूषित कपड़ों और शरीर के अंगों को अच्छी तरह धोएं।
5. छिड़काव के समय धूम्रपान, खाना, पीना और कुछ चबाना नहीं चाहिए।
6. छिड़काव करते या प्रयोग समय पूर्ण सुरक्षात्मक कपड़े पहनें।

विष नाशक

कोई विशिष्ट विषनाशक नहीं हैं। लक्षणानुसार इलाज करें।

प्राथमिक चिकित्सा

कृप्या संलग्न लीफलेट देखें।

लक्षण

कृप्या संलग्न लीफलेट देखें।

उपयोग

एसिटामिप्रिड 20 प्रति. घुलनशील पाउडर कपास के माहू/तेला और सफेद मक्खी; बंदगोभी में माहू; भिन्डी में माहू को रोकथाम के लिये प्रयोग किया जाता हैं ।

19/D, Raj Industrial Park, Karchiya, Savli, Vadodara -
391310 (Gujarat)

निर्माता

अंग्रेजी में लिखे अनुसार

आयात कर्ता का उत्पादन परिसर

अंग्रेजी में लिखे अनुसार

Safety Pictograms / सुरक्षा चित्रलेख



Signature valid

Digitally signed by Archana Sinha
Date: 2025.02.28 16:17:19
Location: CIB&RC-NH4, CGO
Complex, Faridabad,
Designation: Secretary (CIB&RC)

455755-F/9(4)2024

Caution: "Not to Be Used on Crops Other Than Specified on This Label/Leaflet".

Leaflet Details

CIR-346880/2025-Acetamiprid (SP) (461)-1443
Acetamiprid 20% SP
(Insecticide)

Acetamiprid 20% SP is a soluble powder formulation containing 20% of the active ingredient Acetamiprid. Acetamiprid 20% SP is a highly effective systemic insecticide for the control of aphids, jassids and whiteflies in cotton crop; Aphid in Cabbage; Aphid in Okra.

Recommendation

Crop(s)	Common Name of Pest	Dosage/HA		Dilution in Water in Water (liter)	Waiting Period between last spray to harvest days	Re-entry after each Application (In Hours)
		AI (gm)	Formulation (gm)			
Cabbage	Aphid	15.00	75.00	500-600	07	-
	Aphids	15	75	500-600	7	
	Aphids	15	75	500-600	7	
Chilli	Thrips	10-20	50-100	500-600	3	
Cotton	Whiteflies	20	100	500-600	15	
	Jassids	10	50	500-600	15	
	Aphids	10	50	500-600	15	
Okra	Aphids	15	75	500-600	3	
Okra (Bhindi)	Aphid	15.00	75.00	500-600	03	-
Rice	Brown plant hopper	10-20	50-100	500-600	7	

Direction Of Use

Knapsack sprayer, foot sprayer, compression knapsack sprayer, compression knapsack battery sprayer and ASPEE-HTP power sprayer.

Time of Application

Precautions

1. Keep away from foodstuffs, empty foodstuff containers and animals food.
2. Avoid contact with mouth, eyes and skin.
3. Avoid inhalation the spray mist. Spray in the direction of wind.
- 4 . Wash thoroughly the contaminated clothes and parts of the body after spraying.
5. Do not smoke, drink, eat and chew anything while spraying.
6. Wear full protective clothing while mixing and spraying.

Symptoms Of Poisoning

Nausea, vomiting, headache, dizziness and stomach-ache may occur.

First Aid

1. If swallowed, induce vomiting by tickling the back of throat. Repeat it until the vomitus is clear. Do not induce vomiting if the patient is unconscious.
2. If clothing and skin are contaminated remove the clothes and wash the contaminated skin with copious amount of soap and water.
- 3 If eyes are contaminated, flush with plenty of saline/clean water for about 10 to 15 minutes.
4. If inhaled, remove the patient to fresh air.

Phytotoxicity

The product is not phytotoxic when used as per field recommendation.

Antidote

No specific antidote is known. Treat symptomatically.

Disposal Of Used Container

1. Empty containers shall be broken and buried away from habitation
2. The used containers or package shall not be left outside to prevent their re-use
2. The used containers or package shall not be left outside to prevent their re-use
3. The package or surplus material and washing should be disposed off in a safe manner so as to prevent environmental and water pollution

Storage Conditions

1. The packages containing the insecticides shall be stored in separate rooms or premises away from the rooms or premises used for storing other articles or shall be kept in separate almirah under lock and key depending upon the quantity and nature of the insecticide
2. The rooms or premises meant for storing the insecticides shall be well built, dry, well-lit, ventilated and of sufficient dimensions to avoid contamination with vapours

Chemical Composition

S.No	Ingredient	Description	Weight
1	Acetamiprid a.i.		20.00 % w/w
2	Colouring agent	(Brilliant Blue)	0.05 % w/w
3	Sodium Sulphate & Sodium alkylate Mixture		% w/w
4	Blend of Sodium Dodecyl Benzene Sulfonate		4.50 %
5	Sodium Citrate, Dihydrate		30.00 % w/w
6	Calcium Hydrogen Phosphate Dihydrate		0.10 % w/w
7	Lactose Monohydrate		Q.S %
Total			100 % w/w

Manufactured By

M/s SAHAJ AGROCHEM INDUSTRIES

SF 37, Earth Alpha, Vasna Road, Vadodara- 390007 (Gujarat)

Manufacturer Premises Address

19/D, Raj Industrial Park, Karchiya, Savli, Vadodara - 391310 (Gujarat)

पत्रक विवरण

CIR-346880/2025-Acetamiprid (SP) (461)-1443

ऐसिटामिप्रिड 20 प्रतिशत एस पी

(कीटनाशक)

ऐसिटामिप्रिड 20 प्रति. घुलनशील पाउडर कपास के माहू/तेला और सफेद मक्खी; बंदगोभी में माहू; भिन्डी में माहू को रोकथाम के लिये प्रयोग किया जाता हैं ।

उपयोग

फसल	कीट के नाम	प्रति हैक्टेयर मात्रा		पानी की मात्रा (लीटर)	अन्तिम छिड़काव तथा फसल काटने के बीच अन्तराल (दिन)	पुनः प्रवेश की अवधि प्रत्येक छिड़काव के बाद (घंटे)
		स. तत्व (ग्राम)	संरचना (ग्राम)			
बंदगोभी	माहू	15.00	75.00	500-600	07	-
	चेंपा या माहू	15	75	500-600	7	
		15	75	500-600	7	
मिर्च	चुरदा	10-20	50-100	500-600	3	
कपास	सफेद मक्खी	20	100	500-600	15	
	हरा तेला	10	50	500-600	15	
	चेपा या माहू	10	50	500-600	15	
भिन्डी	चेंपा या माहू	15	75	500-600	3	
भिन्डी	माहू	15.00	75.00	500-600	03	-
धान	भुरा फुदका	10-20	50-100	500-600	7	

उपयोग के लिए दिशा-निर्देश

उपयोग में आने वाले यंत्र :नेपसेक स्प्रेयर, फूट स्प्रेयर, कम्प्रेसन नेपसेक बैटरी स्प्रेयर और एस.पी.ई.ई.-एच.टी.पी. पावर स्प्रेयर आदि ।

प्रयोग का समय

प्रयोगकर्ताओं के लिए सावधानियां

1. खाद्य सामग्री , खाद्य सामग्री के खाली बर्तन और पशुओं के चारे से दूर रखें।
2. मुंह, त्वचा और आंखों के सम्पर्क से बचायें।

3. छिडकाव की वाष्प को सांस द्वारा अन्दर जाने से बचायें। हवा की दिशा में छिडकाव करें।
4. छिडकाव के बाद दूषित कपड़ों और शरीर के अंगों को अच्छी तरह धोएं।
5. छिडकाव के समय धूमपान, खाना, पीना और कुछ चबाना नहीं चाहिए।
6. छिडकाव करते या मिलाते समय पूर्ण सुरक्षात्मक कपड़े पहनें।

विष के लक्षण

मितली, उल्टी सिर-दर्द, चक्कर आना, पेट में दर्द हो सकते हैं।

प्राथमिक चिकित्सा

1. यदि निगल जाये तो गले के पीछे गुदगुदी करके उल्टी करायें यह क्रिया तब तक दोहराते रहे जब तक उल्टी द्वारा निकला पदार्थ साफ न हो जाए। यदि मरीज बेहोश हो तो उल्टी न करायें।
2. यदि कपड़े और त्वचा दूषित हो जाए तो दूषित कपड़े उतार दे और त्वचा को काफी मात्रा में साबुन और पानी से धोएं।
3. यदि आंखें दूषित हो जाये तो उनको काफी मात्रा में सैलाइन/साफ पानी से लगभग 10-15 मिनट तक धोएं।
4. यदि खरपतवारनाशक सांस द्वारा अन्दर गया हो तो रोगी को शुद्ध हवा में ले जायें।

पौधविशाक्तता

सुझाव की मात्रा के अनुसार प्रयोग करने पर यह कीटनाशक पौधविशाक्त नहीं हैं ।

विष नाशक

कोई विशिष्ट विषनाशक नहीं हैं। लक्षणानुसार इलाज करें।

खाली डिब्बों का निपटारा

1. कीटनाशक निर्माताओं या उपयोगकर्ता का कर्तव्य है कि डिब्बों या पात्रों को तोड़फोड़कर आबादी से दूर जमीन में गाड़ देना चाहिये।
2. प्रयोग किये गये डिब्बों को दुबारा प्रयोग के लिये बाहर खुला न छोड़ें।
3. डिब्बों या बची हुई कीटनाशक या धोवन का निपटारा इस प्रकार करें कि जिससे आसपास की चीजें, पीने का पानी विषैला न हों

संग्रहण की शर्तें

1. कीटनाशक के डिब्बों को अलग-अलग कमरों या आलमारियों से जिनमें अन्य वस्तुएं विशेषकर खाद्य पदार्थ या पशु आहार संग्रहित किए जाते हैं से दूर अन्य कमरों से दूर अलग-अलग कमरों में या आलमारियों में ताला चाबी में बंद रखें।
2. जिन कमरे अथवा आलमारियों में कीटनाशक का भंडारण करना हो वह अच्छी तरह बना हुआ, सूखा, ठंडा, प्रकाश युक्त,

हवादार, व पर्याप्त लम्बा चौड़ा होना चाहये ताकि कीटनाशक भाप से वातावरण प्रदूषण होने का डर न हों।

रासायनिक संरचना

क्रमांक	घटक	विवरण	वज़न
1	ऐसिटामिप्रिड सक्रि. तत्व		20.00 % भार/ भार
2	रंग लाने वाले घटक	(ब्रिलियन्ट ब्लू)	0.05 % भार/ भार
3	सोडियम सल्फनेट एण्ड सोडियम एल्केलाईट		%
4	सोडियम डोडिसाइल बेन्ज़ीन सल्फोनेट		4.50 % भार/ भार
5	सोडियम साइट्रेट, डाईहाइड्रेट		30.00 % भार/ भार
6	कैल्शियम हाइड्रोजन फॉस्फेट डाईहाइड्रेट		0.10 % भार/ भार
7	लैक्टोज मोनोहाइड्रेट		पर्याप्त मात्रा %
कुल			100 % w/w

निर्माता


अंग्रेजी मे लिखे अनुसार

उत्पादन परिसर

अंग्रेजी मे लिखे अनुसार

Signature valid

Digitally signed by Archana Sinha
Date: 2025.02.28 16:17:21
Location: CIB&RC,NH4,CGO
Complex,Faridabad.
Designation: Secretary (CIB&RC)



455755-F/9(4)2024

Caution:“Not to Be Used on Crops Other Than Specified on This Label/Leaflet”.